



Research Notes

Genetic analysis of the hybrids, involving monostem / shy branching genotypes in sesame (*Sesamum indicum* L.)

P.Sumathi and V. Muralidharan

Abstract

An experiment was conducted with thirty hybrids produced by line x tester mating design from eleven sesame genotypes involving five branched and six monostem / shy branching types. Observations were recorded on days to 50% flowering, days to maturity, plant height, number of branches per plant, number of capsules per plant, capsule length, capsule breadth, number of seeds per plant, 100 seed weight, seed yield per plant and oil content. The traits, number of branches per plant, number of capsules per plant and seed yield per plant showed high PCV and GCV estimates. There is enough scope for selection based on these characters, and the diverse genotypes can provide materials for a sound breeding programme. High heritability combined with high genetic advance (as per cent of mean) observed for plant height, number of branches, number of capsules and seed yield per plant showed that these characters were controlled by additive gene effects and phenotypic selection for these characters would likely to be effective. The characters, plant height, number of branches per plant, number of capsules per plant, days to 50% flowering, days to maturity and 100 seed weight showed significantly positive correlation with seed yield per plant. The trait capsule breadth showed significantly negative association with seed yield per plant.

Key words: Sesame, variability, co-efficient of variation, heritability and genetic advance

Sesame is one of the important and world's oldest edible oilseed crops. Though sesame occupies a place of prominence among oilseeds, its production has been relatively low as compared to other oilseed crops. The major constraints identified for most of the countries including India are, instability in yield, lack of wider adaptability, drought, non – synchronous maturity etc. The branched varieties need more space per plant and hence maintenance of population is very difficult in the commercial planting. This also poses serious problem in the countries where mechanical harvesting is restored. It is preferable to have unicum / monostem / shybranching lines for high-density population for productivity improvement per unit area. Development of hybrids is one of the best method to increase the yield in sesame. By exploiting hybrid vigour sesame cultivation can be revolutionized. The

success of any crop improvement programme essentially depends on the nature and magnitude of variability present in the crop. The knowledge on nature and magnitude of genetic variability is of immense value for planning effective breeding programme to improve the yield potential of the genotypes. The study of interrelationship of yield components is imperative to enable the selection of sesame genotypes. The present investigation was carried out to gather information on variability, heritability and genetic advance and to design the association between yield and yield components in hybrids and parents including single stemmed genotypes of sesame.

Experimental material for the present study consisted of eleven sesame genotypes involving five branched female parents, CO 1, Paiyur 1, TMV 3, TMV4, TMV5 and six monostem / shybranching, male parents viz., MT 34, Cordebergea, KS 990812, KS99037, KS 990813 and KS 99153 and were crossed in line x tester mating design . The resulting 30 hybrids along with their parents were raised

Centre for Plant Breeding and Genetics,
Tamil Nadu Agricultural University, Coimbatore-641003,
Tamil Nadu , India
Email: sumivetri@yahoo.com

during kharif 2005 at Department of oilseeds, Centre for Plant Breeding and Genetics, TNAU, Coimbatore, in a randomized block design with three replications. Each plot consisted of three rows of 4m length spaced at 45 cm between rows and 30 cm between plants. Normal recommended cultural practices and plant protection measures were followed. Ten competitive plants were randomly selected for recording biometrical observations on days to 50% flowering, days to maturity, plant height, number of branches per plant, number of capsules per plant, capsule length, capsule breadth, number of seeds per plant, 100 seed weight, seed yield per plant and oil content. The data were subjected to statistical analysis. The phenotypic and genotypic coefficient of variability was computed as per Burton, 1952. Heritability in broad sense was computed by the formula suggested by Lush (1940). Genetic advance was worked out as per the formula given by Johnson *et al.*, (1955). Correlation coefficients for yield and yield components were evaluated utilizing the formula suggested by Al-jibouri *et al.*, (1958).

Analysis of variance revealed highly significant differences among genotypes for all the characters except capsule breadth indicating considerable amount of genetic variation present in the material. High magnitude of variation in the experimental material was reflected by high values of mean and range for almost all the characters. The minimum number of branches was observed for the hybrids, Paiyur 1 x Cordebergea, TMV 5 x Cordebergea, TMV 5 x MT 34 and TMV 5 x KS 990812 which is ranged from 4-5 branches. The monostem / shybranching parents showed 0 to 2 number of branches only. The maximum number of branches were shown by the branched type parent CO1.

The estimates of genetic parameters like genotypic coefficient of variation, heritability and genetic advance are presented in table 1. Perusal of the data indicates that the traits plant height, number of capsules per plant and seed yield per plant recorded highest phenotypic and genotypic variation than the other characters studied.

Coefficient of variation: The phenotypic coefficient of variation (PCV) was greater than genotypic coefficient of variation (GCV) for all the characters studied, indicating that the environment had an important role in the expression of these characters. The traits, number of branches per plant, number of capsules per plant and seed yield per plant showed high PCV and GCV estimates. There is enough scope for selection based on these characters, and the diverse genotypes can provide materials for a sound breeding programme. Saravanan *et al.*, (2003) and

Solanki and Deepak Gupta (2003) reported high coefficient of variation for number of branches per plant and number of capsules per plant. High coefficient of variation for number of capsules per plant was also reported by Anitha vasline (2000). Plant height showed moderate PCV and GCV and the other traits recorded low PCV and GCV. Sudhakar *et al.* (2007), and Shadakshari *et al.* (1995) reported low phenotypic and genotypic coefficient of variation for the characters days to fifty per cent flowering, days to maturity and oil content. Low coefficient of variation for number of seeds per capsule was reported by Thangavel *et al.*, (2000).

Heritability : The heritability estimates obtained were high for all the characters studied except capsule breadth, which recorded very low heritability for these hybrids and parents. Estimates of heritability and genetic advance in combination are more important for selection than heritability alone. High heritability combined with high genetic advance (as per cent of mean) observed for plant height, number of branches, number of capsules and seed yield per plant showed that these characters were controlled by additive gene effects and phenotypic selection for these characters would likely to be effective. Similar results were reported by Reddy *et al.* (2001) and Krishnaiah *et al.* (2002). The improvement of the characters days to fifty per cent flowering, days to maturity, number of seeds per capsule and 100 seed weight also possible since these characters had high heritability and moderate genetic advance as per cent of mean. The traits capsule length and oil content showed high heritability with low genetic advance. These results are in conformity with the findings of Reddy *et al.* (2001) and Sudhakar *et al.* (2007). The trait capsule breadth showed low heritability and low genetic advance such situation may arise due to non additive gene action.

Association analysis: Yield is a complex quantitative trait, greatly influenced by environmental fluctuations. Hence, selection based on yield performance alone may indicate a biased result and leads to ambiguity. A study of nature and degree of association of component characters with yield assumes greater importance for fixing up characters that play a decisive role in influencing yield. Selection would therefore be more effective, if it is based on component characters rather than directly on yield. Correlation coefficient analysis measures the mutual relationship between various characters and is used to determine the component character on which selection can be done for improvement in yield.

The characters plant height, number of branches per plant, number of capsules per plant, days to 50% flowering, days to maturity and 100 seed weight showed significantly positive correlation with seed yield per plant (Table 2). The trait capsule breadth showed significantly negative association with seed yield per plant. Similar results were reported by Pawar *et al.* (2002); and Deepa Sankar and Ananda Kumar (2003) for plant height; Ramireddy Kumar and Sundaram (2002) and Deepa Sankar and Ananda Kumar (2003) for number of branches; Ramireddy Kumar and Sundaram (2002), Pawar *et al.* (2002) and Deepa Sankar and Ananda Kumar (2003) for number of capsules per plant.

The trait plant height showed significantly positive correlation with the traits number of branches per plant, number of capsules per plant, days to 50% flowering, days to maturity and 100 seed weight. Earlier reports of Ramireddy kumar and Sundaram (2002) and Deepa Sankar and Ananda kumar (2003) revealed the positive association of plant height with number of branches per plant and number of capsules per plant. The trait, number of branches per plant showed significantly positive inters correlations with number of capsules per plant, days to 50% flowering, days to maturity, 100 seed weight. Reddy *et al.*, (2001) and Pawar *et al.*, (2002) reported the positive association of number of branches per plant with number of capsules per plant. Similarly, the trait, number of capsules per plant recorded positive correlation with 50% flowering, days to maturity, 100 seed weight but it showed negative significant association with capsule breadth. Days to 50 per cent flowering showed positive association with days to maturity and negative association with capsule length, days to maturity revealed positive association with 100 seed weight and number of seeds per capsule showed positive association with capsule length.

The present study indicated that the selection based on the characters, plant height, number of branches per plant, number of capsules per plant, and seed yield per plant would be effective for the development of superior sesame variety through recombination breeding.

References

- Al-Jibouri HA, Miller PA and Robinson HF 1958 Genotypic and environment variances and covariances in an upland cotton cross of interspecific origin. *Agron. J.*, 50 : 633-636.
- Anitha Vasline Y, Saravanan K and Ganesan J 2000 Studies on variability and genetic advance for certain characters in mutant populations of sesame (*Sesamum indicum* L.). *Sesame and Safflower Newsl.*, 15 : 39-43.
- Burton GW 1952 Quantitative inheritance in grasses. *Proc. Sixth Int. Grassland Congr.* 1: 277-283.
- Deepa Sankar P and Ananda Kumar CR 2003 Genetic analysis of yield and related components in sesame (*Sesamum indicum* L.) *Crop Res.* 25(1) : 91-95.
- Johnson HW Robinson HF and Comstock RE 1955 Estimation of genetic and environmental variability in soybean. *Agron. J.* 47: 314-318.
- Krishnaiah G, Reddy KR and Sekar MR 2002 Variability studies in sesame. *Crop Res.* 24: 3, 501-504.
- Lush JL 1940 Intraclass correlation and regression of offspring on dams as a method of estimating heritability of characters. *Proc. Amer. Soc. Anim. Nutr.* 32: 293-301.
- Pawar KN, Chetti MB and Shamrao Jahagirdar, 2002. Association between seed yield and yield attributing characters in sesame (*Sesamum indicum* L.) *Agri. Sci. Digest*, 22(1) : 18 – 20.
- Ramireddy Kumar A and Sundaram T 2002 Inter relationships among yield and yield components in sesame (*Sesamum indicum* L.) *J. Res. ANGRAU* 30 (2) : 42-44.
- Reddy PAV, Sekar MR, Ranganatha ARG and Dhanraj A 2001 Genetic variability and heritability for seed yield and its components in sesame (*Sesamum indicum* L.). *J. Oilseeds Res.* 18 : 2, 173-175.
- Saravanan S and Nadarajan N 2003. Combining ability studies in sesame. *Crop Res.* 25(2) : 319 – 324.
- Shadakshari YG, Virupakshappa K and Shivashankar G 1995 Genetic variability studies in germplasm collection of sesame (*Sesamum indicum* L.). *Mysore J. Agric. Res.* 29 : 133-137.
- Solanki ZS and Deepak Gupta 2003 Variability and character association among quantitative characters of sesame. *J. Oilseeds Res.* 20 : 276-277.
- Sudhakar N, Sridevi O and Salimath P M 2007 Variability and character association analysis in sesame, *Sesamum indicum* L. *J Oilseeds Res.* 24 : 1, 56-58.
- Thangavel P, Saravanan K, Senthil Kumar P, Anbuslvan, Y and Ganesan J 2000. Variability, heritability and genetic advance in sesame (*Sesamum indicum* L.) *Sesame and Safflower Newsl.*, 15 : 19 – 22.

**Table 1. Estimates of variability parameters for yield and its components**

Character	Mean	Range	Genotypic variance	Phenotypic variance	Genotypic coefficient of variation (%)	Phenotypic coefficient of variation (%)	Heritability %	Genetic advance as percent of mean
Plant height (cm)	121.26	48.6-153.7	413.60	448.63	16.77	17.47	92.2	33.17
No. of primary branches / plant	6.66	0.33-11.87	7.07	8.38	39.94	43.48	84.4	75.56
No. of capsules/ plant	152.85	21.1-224.6	2063.54	2169.35	29.72	30.47	95.12	59.71
Days to 50% flowering	46.28	35.3-52.3	12.01	16.93	7.49	8.89	70.90	12.99
Days to maturity	86.15	74.3-94.3	26.46	27.54	5.97	6.09	96.1	12.06
No. of seeds/ capsule	56.63	40.6-65.2	22.09	32.70	8.93	10.87	67.6	15.12
100 seed weight (g)	0.33	0.29-0.38	0.0005	0.0006	6.81	7.39	84.9	12.91
Capsule length (cm)	2.58	2.33-2.97	0.02	0.02	5.18	5.62	84.9	9.82
Capsule breadth (cm)	0.72	0.67-0.80	0.0009	0.0011	4.15	4.59	81.8	7.74
Oil content (%)	41.59	39.3-43.73	1.38	1.38	2.64	2.83	87.1	5.07
Seed yield/plant(g)	21.04	5.8-30.9	48.23	48.23	31.49	33.00	91.0	61.89

**Table 2. Genotypic correlation coefficients between different characters in sesame**

Characters	No. of branches/ Plant	No. of capsules / plant	Days to 50% flowering	Days to maturity	No. of seeds/ capsule	100 seed weight	Capsule length	Capsule breadth	Oil content	Seed yield / plant
Plant height	0.949**	0.860**	0.872**	0.894**	-0.163	0.632*	-0.323	-0.333	0.407	0.824**
No. of primary branches	1.000	0.833**	0.919**	0.976**	-0.349	0.701**	-0.3628	-0.415	0.241	0.835**
No. of capsules/ plant		1.000	0.741**	0.772**	-0.1855	0.601	-0.2834	-0.786**	0.293	0.92**
Days to 50% flowering			1.000	0.947**	-0.2358	0.429	-0.595*	-0.1900	0.308	0.682**
Days to maturity				1.000	-0.3523	0.547	-0.4153	-0.548*	0.366	0.766**
No. of seeds/ capsule					1.000	-0.204	0.594*	0.2814	0.434	-0.031
100 seed weight						1.000	-0.2392	-0.883**	0.094	0.566**
Capsule length							1.000	0.0074	-0.205	-0.263
Capsule breadth								1.000	-0.477	-0.953**
Oil content									1.000	0.413

*Significant at P = 0.05

** Significant at P = 0.01