

Electronic Journal of Plant Breeding



Research Article

Genome editing of elite rice cultivar CO51 for bacterial leaf blight resistance

T. Arulganesh¹, Yaiphabi Kumam¹, K. K. Kumar¹, L. Arul¹, E. Kokiladevi¹, S. Nakeeran¹, S. Varanavasiappan¹, S. Manonmani² and D. Sudhakar^{1*}

¹Department of Plant Biotechnology, Centre for Plant Molecular Biology and Biotechnology, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India.

² Department of Plant Genetic Resources, Centre for Plant Breeding and Genetics, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India.

*E-Mail: dsudhakar@hotmail.com

Abstract

Bacterial leaf blight (BLB) of rice caused by *Xanthomonas oryzae* pv *oryzae* (Xoo) is a major constraint in rice production. Natural resistance to BLB in some of the wild type cultivars of rice is due to mutations in effector binding element (EBE), present in the promoter region of *SWEET* genes. *SWEET14* is one of the most commonly targeted genes by majority of the TALEs of Xoo pathogen. Thus, the present study was aimed at creating novel mutations in the EBE of *OsSWEET14* gene in *indica* rice cultivar, CO51 through CRISPR/Cas9 mediated genome editing technology to impart resistance against BLB. *Agrobacterium*-mediated transformation using immature embryos followed by regeneration resulted in eleven transgenic plants from six independent transformation events, of which nine plants (belonging to five events) had mutations in the target sequence. Bioassay studies on four mutant plants (belonging to three events) resulted in the identification of two plants (belonging to two events) with resistance/moderately resistance against BLB.

Key words: Rice, *SWEET* genes, CRISPR/Cas9, Bacterial Leaf Blight, Effector Binding Element

INTRODUCTION

Rice (*Oryza sativa*) is a cereal crop that is consumed by half of the global population (Fukagawa and Ziska, 2019). High population growth, limited arable land, shortage of other natural resources and outbreak of crop diseases limit rice cultivation and production (FAO, 2017). Among these constraints, yield loss due to diseases is a serious concern, contributing up to 16 per cent globally (Ficke *et al.*, 2018). Bacterial leaf blight (BLB) is one of the major foliage diseases that attack the rice crop seasonally. Yield losses due to BLB have been reported to vary from 2 to 74 per cent depending on location, season, crop growth stage and cultivar (Rao and Kauffman, 1977; Reddy *et al.*, 1978). Major symptoms of BLB include water soaked yellowish stripes on leaf blades, curling of leaves and early morning appearance of bacterial ooze on

young lesions on the leaves (Nino-Liu *et al.*, 2006). The pathogen, *Xanthomonas oryzae* pv. *oryzae* (Xoo) targets *SWEET* (Sugar will eventually be exported transporters) genes in rice during BLB infection. Even though the rice genome has more than 20 *SWEET* genes, only three (*OsSWEET11*, *OsSWEET13* and *OsSWEET14*) are targeted by the Xoo pathogen, and these genes are called 'susceptible (S) genes' (Oliva *et al.*, 2019). During infection with BLB, the Xoo pathogen secretes Transcription Activator like Effector (TALE) molecules which bind to the effector binding element (EBE) in the promoter region of some of the rice *SWEET* genes. The binding of TALEs to the promoter region results in the over-expression of *SWEET* genes. Over-expression of *SWEET* genes results in the production of sugars,

rendering the plant susceptible to the pathogen (Oliva *et al.*, 2019). There are six known TALE proteins that target *OsSWEET* gene promoter regions, of which four different TALEs viz., PthXo3, AvrXa7, TalC, and TalF (previously known as Tal5), target EBE present in *OsSWEET14* gene (Blanvillain-Baufume *et al.*, 2017). The remaining two TALEs, PthXo1 and PthXo2 target EBE present in *OsSWEET11* and *OsSWEET13* genes respectively (Oliva *et al.*, 2019). Naturally occurring recessive alleles of these *SWEET* genes confer resistance against BLB (Zafar *et al.*, 2020). Recessive alleles, *xa13* and *xa25* for the genes *OsSWEET11* and *OsSWEET13* were found to confer resistance against BLB caused by *Xoo* race 6 (Chu *et al.*, 2006) and *Xoo* strain PXO339 (Chen *et al.*, 2002), respectively. With respect to the *OsSWEET14* gene, *xa41* recessive allele identified in a few of the African wild type cultivars, *Oryza barthii* and *O. glaberrima* conferred resistance to BLB (Hutin *et al.*, 2015). Most of the recessive alleles contain nucleotide polymorphisms in the EBE of *OsSWEET* genes (Oliva *et al.*, 2019).

Genome editing is a novel biotechnological approach that promises to create nucleotide modifications at the target region within a gene. Previous studies have used genome editing technologies such as CRISPR-Cas9 to generate resistance against BLB by creating targeted mutations in the EBE of *OsSWEET14* genes (Oliva *et al.*, 2019; Blanvillain-Baufume *et al.*, 2017; Zafar *et al.*, 2020). Though there are several types of CRISPR systems available, the Cas9 nuclease from *Streptococcus pyogenes* (a type II CRISPR/Cas9 system) is most commonly used (Li *et al.*, 2020b). Cas9 creates a double stranded break at the target region based on PAM (proto-spacer adjacent motif) sequence 'NGG' which is later repaired either by non-homologous end joining (NHEJ) or homology directed repair (HDR) mechanism (Jiang and Doudna, 2017). Since the CRISPR/cas9 induced mutation occurs in the T-DNA transformed rice calli before the regeneration stage, chances are there to identify homozygous mutants

with high mutation frequencies in the T_0 generation itself. At the same time, mutants without exogenous T-DNA can be segregated out in the T_1 generation. These mutants are also known as "transgene-free" mutants. (Ricroch *et al.*, 2017). Considering these available opportunities, the present study was aimed towards the development of BLB resistant lines in the genetic background of a short duration *indica* rice cultivar CO51, by using CRISPR/Cas9 technology.

MATERIALS AND METHODS

Plants of the high yielding, popular short duration rice genotype, CO51 were maintained at Paddy Breeding Station, Tamil Nadu Agricultural University, Coimbatore for regular supply of immature seeds. The immature seeds were collected from the field and used as the source of explants.

The nucleotide sequence of promoter region encompassing the EBE of *OsSWEET14* gene (*Os11g0508600*) was retrieved from the online database tool, EnsemblPlants (<https://plants.ensembl.org/index.html>). This sequence information was used to design single guide RNA (sgRNA) using the web-based tool, CRISPR P 2.0 (<http://crispr.hzau.edu.cn/>) (Fig. 1). The design of sgRNA was restricted to the 29 bp EBE sequence of the promoter region. This identified a promising sgRNA sequence with top strand: 5' ATATAAACCCCTCCAACC 3' and bottom strand 5' GGTTGGAGGGGTTTATAT 3'. During the synthesis of these DNA oligomers, *BsaI* adaptors (5'GGCA 3' for the top strand and 5' AAAC 3' for the bottom strand) were added to facilitate cloning in the *BsaI* restriction site of the binary vector, pRGE32. The plasmid, pRGE32, was a gift from Dr Yinong Yang's lab (Addgene plasmid # 63142; <http://n2t.net/addgene:63142>; RRID: Addgene_63142) (Xie *et al.*, 2015). The T-DNA of the binary vector harbouring the sgRNA sequence is represented in Fig. 2.




Fig. 1. Genomic structure of *OsSWEET14* gene containing effector binding elements (EBEs) for the Transcription-Activator-Like Effectors (TALE) proteins AvrXa7 and PthXo3. The 29 bp EBE is underlined. The sgRNA sequence is in red and the protospacer adjacent motif PAM sequence (AGG) is highlighted in blue.




Fig. 2. T-DNA of binary vector pRGEB32 harbouring the sgRNA of *OsSWEET14* gene

The recombinant binary vector, pRGEB32 harbouring the *Cas9/OsSWEET14sgRNA* was mobilized into *Agrobacterium* strain, LBA4404 by triparental mating. Immature embryos of CO51 were collected from maturing panicles (12-14 days after anthesis). *Agrobacterium*-mediated transformation was done using these immature embryos, following the protocol given by Hiei and Komari (2008) with minor modifications (Nithya *et al.*, 2020; Shweta *et al.*, 2020). Friable, well proliferated yellow calli that survived two rounds of stringent selection in 50 mg^l⁻¹ hygromycin were successfully regenerated into complete plants on regeneration medium (RNMH30 medium; Hiei and Komari, 2008) and rooted on half strength MS medium (Murashige and Skoog, 1962). Finally, the plants were hardened in pro-trays and maintained in the transgenic greenhouse. The transformation efficiency (%) was calculated by using the following formula,

$$\frac{\text{Number of cocultivated embryos that produced plants}}{\text{Total number of embryos cocultivated}} \times 100$$

To determine the mutations at the target site, molecular characterization of putative T₀ mutants was done. Genomic DNA was isolated from wild type and T₀ mutant plants using the CTAB method (Porebski *et al.*, 1997). PCR analyses for the presence of *cas9* and *hpt* genes in these mutants were performed using sequence specific primers with respective temperature profiles (Table 1). Further, the target region of about 344 bp encompassing the sgRNA sequence was also amplified using sequence specific primers (Table 1). The amplified products were purified and sequenced by the Sanger method (Eurofins, Bengaluru). The sequence results were analyzed using the online software, DS DecodeM (<http://skl.scau.edu.cn/dsdecode/>) (Xie *et al.*, 2017; Liu *et al.*, 2015) and CRISP I.D. (<http://crispid.gbiomed.kuleuven.be/>) (Dehairs *et al.*, 2016) to detect the mutants by comparing the sequence of transgenic plants with those of wild type plant. The mutation efficiency was calculated by using the following formula,

$$\text{Mutation efficiency} = \frac{\text{Number of events with mutations}}{\text{Total number of events generated}} \times 100$$

Bioassay on T₀ rice mutant plants for resistance against

BLB was performed in the transgenic greenhouse. The leaf clipping method proposed by Kauffman *et al.* (1973) was followed for bioassay of T₀ mutant plants against *Xoo* strain, 1X020 which was obtained from ICAR-Indian Institute of Rice Research, Hyderabad. For the leaf clipping method, a three-day-old culture of *Xoo* maintained on nutrient agar medium was used. Fresh *Xoo* suspension was prepared by suspending a loopful of culture in 10 Mm sterile MgCl₂. The optical density of the bacterial suspension was adjusted to 0.5 at 600 nm using a spectrophotometer (Yinggen *et al.*, 2017). Forty five day-old plants belonging to four mutant plants (from three events), a transformed (wild type) plant and a non-transformed CO51 (wild type) plant were subjected to bioassay. For this, sterilized scissors (in 70 % ethanol) were dipped in *Xoo* suspension and used to cut about 4-5 cm of first three fully extended leaves of T₀ adult plants as well as wild type plants. Additionally, sterilized scissors dipped in MgCl₂ without *Xoo* suspension were also used to cut the leaves and it served as the negative control. Observations on symptom development in the plants were recorded 14 days post inoculation (dpi). Lesion length on the inoculated leaves was measured and recorded. The data obtained were subjected to Tukey's HSD test using the online statistic tool OPSTAT (Sheoran *et al.*, 1998) to identify any significant difference between the mutant and wild type plants.

RESULTS AND DISCUSSION

Within a decade of the development, CRISPR/Cas9 based genome editing technology has emerged as a revolutionary genetic engineering technology accelerating crop improvement programs. The edited plants developed are a result of small indels or substitutions of nucleotides within the endogenous target gene, resulting in novel alleles. Using CRISPR tools, several novel alleles for grain number, grain size, panicle structure (Li *et al.*, 2016) disease resistance (Li *et al.*, 2020a; Oliva *et al.*, 2019), herbicide tolerance (Sun *et al.*, 2016), aroma (Ashokkumar *et al.*, 2020), grain quality (Huang *et al.*, 2020), nutritional quality (Zafar *et al.*, 2020) and herbicide tolerance (Sun *et al.*, 2016) have been generated in rice.

Various approaches have been used in the past to develop disease resistant rice cultivars, which include the introgression of resistance genes (R genes), host-derived resistance, and loss of susceptibility through mutation

Table 1. List of primers used in this study

Name of gene	Primer sequence (5' to 3')	Amplicon size (bp)	PCR conditions
<i>hpt*</i>	Forward primer: GCTGTTATGCGGCCATTGGTC Reverse primer: GCCTCCAGAAGAAGATGTTG	686	94 °C for 5 minutes 94 °C for 45 seconds 57.8 °C for 45 seconds 72 °C for 45 seconds 72 °C for 10 minutes ×30 cycles
<i>cas9*</i>	Forward primer: CTTCTGGCGGTTCTCTTAG Reverse primer: TGCTGTTTGATCCGTTGTTG	478	95 °C for 5 minutes 95 °C for 45 seconds 52 °C for 45 seconds 72 °C for 45 seconds 72 °C for 10 minutes ×30 cycles
<i>SWEET14</i>	Forward primer: CATGGCTGTGATTGATCAGG Reverse primer: GTTTGGTGGGAGGAGATCAA	344	94 °C for 5 minutes 94 °C for 45 seconds 54 °C for 45 seconds 72 °C for 45 seconds 72 °C for 10 minutes 4 °C for infinite hold ×30 cycles

* Nithya *et al.* (2020)

in recessive R genes. Even though the introgression of R genes is very effective against *Xoo* pathogens, the large scale deployment of the R gene may result in the emergence of new virulent *Xoo* strains (Ji *et al.*, 2016). So there is a necessity for alternative approaches which include biotechnological tools such as genome editing for imparting resistance against bacterial blight disease in rice.

One of the BLB susceptible genes in rice is *SWEET14* and mutations in its promoter region, particularly in EBE, resulted in BLB resistance in rice (Oliva *et al.*, 2019; Duy *et al.*, 2021). A sgRNA was designed based on the EBE of *SWEET14* and cloned in a binary vector, pRGEB32, containing sgRNA expression cassette, *cas9* gene and *hpt* gene. The recombinant pRGEB32 was mobilized into *Agrobacterium* strain, LBA4404 and the resultant *Agrobacterium* was used in transformation experiments.

Immature embryos (232 embryos) collected from rice variety CO51 were co-cultivated with the *Agrobacterium* strain harbouring recombinant pRGEB32 possessing *cas9* and *SWEET14* sgRNA expression cassettes. Calli derived from 70 co-cultivated embryos survived two rounds of hygromycin selection and they were subcultured onto regeneration medium for shoot development.

Eleven plants belonging to six independent events were generated with a transformation efficiency of 2.6% (Table 2). Each of the transformation events had two plants except the event TAG-CO51-3 which had only one plant (Table 3).

Molecular analyses by PCR for the presence of *cas9* and *hpt* genes confirmed their presence in all the putative mutants generated (Fig. 3 and Fig. 4). Analysis of Sanger sequencing after amplification of the 344 bp target region identified mutants in five events, thereby resulting in high mutation efficiency of 83.3 % (Table 2). These five events had 9 plants, of which five had a mutation in biallelic, one in monoallelic (heterozygous) and three in biallelic-homozygous forms (Table 3). Five out of the nine mutant plants had mutations at 3 bp upstream of the PAM sequence (Table 3). This indicated the specificity of the CRISPR/Cas9 editing system in the rice genome.

Four mutant plants, one transformed plant (but turned as wild type due to absence of mutation) and CO51 wild type (non-transformed) were subjected to bioassay (Table 4). Other mutant plants could not be included in this study as they were relatively older than mutants tested. Symptom development of BLB infection was observed at 14 days post inoculation. The infection among the mutants and

Table 2. Transformation and mutation efficiency of *Agrobacterium*-mediated transformation of rice cultivar CO51

No. of immature embryos co-cultivated	No. of calli survived two rounds of selection	No. of events generated	Events with mutation	Transformation efficiency (%)	Mutation efficiency (%)
232	70*	6	5	2.5	83.3

* Calli derived from 70 co-cultivated embryos survived two rounds of selection.




Fig. 3. PCR analysis for the presence of *cas9* gene in putative T_0 mutants




Fig. 4. PCR analysis for the presence of *hpt* gene in putative T_0 mutants

Table 3. Details of nucleotide sequence of the target region in CO51 mutants (T₀)

Event I.D	Plant I.D. number	Nucleotide sequence	zygosity
	CO51	WT:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC	Wild type
TAG-CO51-1	TAG-CO51-1-1	A1:TCCTTCCTAGCACTATATAAACCCCTCCATACCAGGTGCTAAGCTC A2:TCCTTCCTAGCACTATATCCACCCCTCCATACCAGGTGCTAAGCTC	Biallelic (1i/2s1i)
	TAG-CO51-1-2	A1:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC A2:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC	Homozygous (1i/1i)
TAG-CO51-2	TAG-CO51-2-1	A1:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC A2:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC	Wild type
	TAG-CO51-2-2	A1:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC A2:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC	Wild type
TAG-CO51-3	TAG-CO51-3-1	A1:TCCTTCCTAGCACTATATAAACCCCTCCA-CCAGGTGCTAAGCTC A2:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC	Biallelic (1d/1i)
TAG-CO51-4	TAG-CO51-4-1	A1:TCCTTCCTAGCACTATATAAACCCCTCCA-CCAGGTGCTAAGCTC A2:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC	Biallelic (1d/1i)
	TAG-CO51-4-2	A1:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC A2:TCCTTCCTAGCACTATATAAACCCCTCCA-CCAGGTGCTAAGCTC	Monoallelic (heterozygous) (WT/2d)
TAG-CO51-5	TAG-CO51-5-1	A1:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC A2:TCCTTCCTAGCACTATATAAACCCCTCCATACCAGGTGCTAAGCTC	Biallelic (1i/1i)
	TAG-CO51-5-2	A1:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC A2:TCCTTCCTAGCACTATATAAACCCCTCCATACCAGGTGCTAAGCTC	Biallelic (1i/1i)
TAG-CO51-6	TAG-CO51-6-1	A1:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC A2:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC	Homozygous (1i/1i)
	TAG-CO51-6-2	A1:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC A2:TCCTTCCTAGCACTATATAAACCCCTCCAACCAGGTGCTAAGCTC	Homozygous (1i/1i)

'-' denotes deletion. Insertions are in red. PAM (AGG) sequence is highlighted in blue and EBE sequence in the wild type is underlined.

wild type plants was compared in terms of lesion length covered by the pathogen in the inoculated leaf. All the inoculated leaves of both wild type and mutant plants expressed typical water soaked lesions of BLB. Among them, event TAG-CO51-5-1 showed the lowest lesion length of 6.5 cm (Fig. 5). Mean comparisons of the lesion lengths using Tukey's HSD test further indicated that it was significantly different from the wild type CO51 which had a lesion length of 15.5 cm. In contrast, no significant difference was observed in the remaining events (Table 4). No lesion was observed in the leaves of uninoculated (cut with scissors without Xoo) wild type plants. The disease resistance scorecard proposed by Duy *et al.* (2021) was used in this study. The events TAG-CO51-5-1 and TAG-CO51-4-1 with lesion lengths of 6.5 cm and 11 cm were classified as resistant (lesion length < 8.0cm) and moderately resistant (lesion length 8-12 cm) against BLB respectively. Both these mutants were biallelic. The only mono-allelic (heterozygous) mutant TAG-CO51-4-2 with a lesion length of 19.33 cm was highly susceptible to BLB, as it has an intact EBE in one of its alleles. This functional allele would allow Xoo to bind its TALE protein and cause the disease. Identification of only two resistant lines in the present study

may be attributed to earlier reports suggesting that larger nucleotide polymorphisms need to be created in the EBE region (Oliva *et al.*, 2019). All the mutants generated in the study had deletions or insertions restricting to only 1 bp. This highlights the need for not only developing larger nucleotide polymorphisms but also screening a larger T₀ mutant population so that more plants with desired mutations could be identified.

Three of the events had mutations in homozygous condition, indicating that CRISPR/Cas9 system induces mutation in the embryonic cell even before it undergoes the first cell division. Besides, all the three homozygous T₀ plants had 1 bp insertion which is in agreement with the earlier report of Zhang *et al.* (2014). Such precise, desirable, targeted mutation in a short time frame demonstrates the high mutation efficiency of the CRISPR/Cas9 system in rice. Besides, this tool greatly saves the time taken to develop a product compared to traditional mutational breeding. In our study, the mutation efficiency of CRISPR/Cas9 for targeting the OsSWEET14 promoter was 83.3 per cent, which was higher than the mutation efficiency of TALENS, which was 51 per cent for the same target sequence (Blainvillain-




Fig. 5. Bacterial leaf blight lesion lengths at 14 days post inoculation in T_0 mutants

Table 4. Lesion length in T_0 mutants of CO51 and wild type CO51

Treatment (Mutations in the T_0 events)	Mutation / zygosity	Mean lesion length (cm) * (Mean \pm S.E)	Resistance reaction
Non-transformed Wild Type CO51	Wild type	15.50 \pm 1.803 ^a	S
TAG-CO51-2-1	Wild type	15.16 \pm 1.691 ^a	S
TAG-CO51-4-1	Biallelic (1d/1i)	11.00 \pm 0.764 ^{ab}	MR
TAG-CO51-4-2	Monoallelic (WT/2d)	19.33 \pm 2.048 ^a	S
TAG-CO51-5-1	Biallelic (1i/1i)	6.50 \pm 2.566 ^b	R
TAG-CO51-6-1	Homozygous (1i/1i)	15.33 \pm 1.424 ^a	S

*Significant difference was calculated at C.D value 5.5346.

Data represent mean \pm S.E of lesion length of three independent replications at 14 dpi Xoo strain, IX020.

Means followed by the same superscript do not differ significantly at $P < 0.05$ (one-way ANOVA with Tukey's HSD test)

baufume *et al.*, 2017). This shows the superiority of CRISPR/Cas9 over other genome editing tools. Moreover, previous reports suggested that RNA mediated silencing of *OsSWEET14* in BLB resistant Kitaake lines caused a negative impact on seed filling and seed production (Antony *et al.*, 2010). In contrast, CRISPR/Cas9 generated mutants of the *OsSWEET14* gene did not show significant differences in major agronomic traits compared to the respective wild type (Duy *et al.*, 2021). So, it is conceivable that minor changes in promoter regions using CRISPR/Cas9 have no effect on the expression of *SWEET* genes as well as the growth and development of plants.

In conclusion, the two BLB resistant / moderately resistant

mutants, TAG-CO51-5-1 and TAG-CO51-4-1 identified from the present investigation, with mutations in the EBE region and reduced lesion length can be promising candidates. Since there is a chance of chimeric mutation and both of the resistant lines are biallelic, their resistance against BLB, however, needs to be proven by bioassay on homozygous progeny population. Besides, their agronomic performances also need to be characterized, although the majority of previous studies suggested that CRISPR/Cas9 induced mutations in the *OsSWEET14* promoter did not negatively impact the main agronomic traits (Duy *et al.*, 2021). Taken together, our findings indicate that targeted modifications of susceptibility (S) *OsSWEET14* gene via CRISPR/Cas9 is effective in imparting resistance against BLB in rice.

ACKNOWLEDGEMENT

The authors thank Dr G.S. Laha, Principal Scientist, Department of Plant Pathology, ICAR-IIRR, Hyderabad for providing the *Xanthomonas* strains for conducting bioassay. The first author expresses his sincere gratitude to DBT for providing the fellowship.

REFERENCES

- Antony, G., Zhou, J., Huang, S., Li, T., Liu, B., White, F. and Yang, B. 2010. Rice *xa13* recessive resistance to bacterial blight is defeated by induction of the disease susceptibility gene *Os-11N3*. *Plant Cell*, **22**: 3864-76. [\[Cross Ref\]](#)
- Ashokkumar, S., Jaganathan, D., Ramanathan, V., Rahman, H., Palaniswamy, R., Kambale, R. and Muthurajan, R. 2020. Creation of novel alleles of fragrance gene *OsBADH2* in rice through CRISPR/Cas9 mediated gene editing. *PLoS One*, **15**: e0237018. [\[Cross Ref\]](#)
- Blanvillain-Baufume, S., Reschke, M., Sole, M., Auguy, F., Doucoure, H., Szurek, B., Meynard, D., Portefaix, M., Cunnac, S., Guiderdoni, E. and Boch, J. 2017. Targeted promoter editing for rice resistance to *Xanthomonas oryzae* pv. *oryzae* reveals differential activities for *SWEET 14* inducing TAL effectors. *Plant Biotechnol. J.*, **15**: 306–317. [\[Cross Ref\]](#)
- Chen, H., Wang, S. and Zhang, Q. 2002. New gene for bacterial blight resistance in rice located on chromosome 12 identified from Minghui 63, an elite restorer line. *Phytopathology*, **92**: 750–754. [\[Cross Ref\]](#)
- Chu, Z., Yuan, M., Yao, J., Ge, X., Yuan, B., Xu, C., Li, X., Fu, B., Li, Z., Bennetzen, J.L. and Zhang, Q. 2006. Promoter mutations of an essential gene for pollen development result in disease resistance in rice. *Genes Dev.*, **20**: 1250–1255. [\[Cross Ref\]](#)
- Dehairs, J., Talebi, A., Cherifi, Y. and Swinnen, J.V. 2016. CRISP-ID: decoding CRISPR mediated indels by Sanger sequencing. *Sci. Rep.*, **6**: 28973. [\[Cross Ref\]](#)
- Duy, P.N., Lan, D.T., Pham Thu, H., Thi Thu, H.P., Nguyen Thanh, H., Pham, N.P. and Pham, X. H. 2021. Improved bacterial leaf blight disease resistance in the major elite Vietnamese rice cultivar TBR225 via editing of the *OsSWEET14* promoter. *PLoS One*, **16**: 255470. [\[Cross Ref\]](#)
- FAO. 2017. *The future of food and agriculture – Trends and challenges*. Rome.
- Ficke, A., Cowger, C., Bergstrom, G. and Brodal, G. 2018. Understanding yield loss and pathogen biology to improve disease management: Septoria Nodorum Blotch - A Case Study in Wheat. *Plant Dis.*, **102**: 696-707. [\[Cross Ref\]](#)
- Fukagawa, N.K. and Ziska, L.H. 2019. Rice: Importance for global nutrition. *J. Nutr. Sci. Vitaminol.*, **65**: S2-S3. [\[Cross Ref\]](#)
- Hiei, Y. and Komari, T. 2008. *Agrobacterium*-mediated transformation of rice using immature embryos or calli induced from mature seed. *Nat. Protoc.*, **3**: 824-34. [\[Cross Ref\]](#)
- Huang, L., Li, Q., Zhang, C., Chu, R., Gu, Z., Tan, H. and Liu, Q. 2020. Creating novel *Wx* alleles with fine-tuned amylose levels and improved grain quality in rice by promoter editing using CRISPR/Cas9 system. *Plant Biotechnol. J.*, **18**: 2164. [\[Cross Ref\]](#)
- Hutin, M., Sabot, F., Ghesquière, A., Koebnik, R. and Szurek, B. 2015. A knowledge-based molecular screen uncovers a broad-spectrum *OsSWEET14* resistance allele to bacterial blight from wild rice. *Plant J.*, **84**: 694-703. [\[Cross Ref\]](#)
- Ji, Z., Ji, C., Liu, B., Zou, L., Chen, G. and Yang, B. 2016. Interfering TAL effectors of *Xanthomonas oryzae* neutralize R-gene-mediated plant disease resistance. *Nat. Commun.*, **7**: 1-9. [\[Cross Ref\]](#)
- Jiang, F. and Doudna, J.A. 2017. CRISPR-Cas9 Structures and Mechanisms. *Annu. Rev. Biophys.*, **22**: 505-529. [\[Cross Ref\]](#)
- Kauffman, H.E., Reddy, A.P.K., Hsieh, S.P.Y. and Merca, S.D. 1973. An improved technique for evaluating resistance of rice varieties to *Xanthomonas oryzae*. *Plant Diseases Reporter*, **57**: 537-541.
- Li, C., W. Li, Z. Zhou, H. Chen, C. Xie, and Y. Lin. 2020a. A new rice breeding method: CRISPR/Cas9 system editing of the *Xa13* promoter to cultivate transgene-free bacterial blight-resistant rice. *Plant Biotechnol. J.*, **18**: 313-315. [\[Cross Ref\]](#)
- Li, H., Yang, Y., Hong, W., Huang, M., Wu, M. and Zhao, X. 2020b. Applications of genome editing technology in the targeted therapy of human diseases: mechanisms, advances and prospects. *Sig Transduct. Target Ther.*, **5**: 1. [\[Cross Ref\]](#)
- Liu, W., Xie, X., Ma, X., Li, J., Chen, J. and Liu, Y.G. 2015. DSDcode: A web-based tool for decoding of sequencing chromatograms for genotyping of targeted mutations. *Mol. Plant.*, **8**: 1431-1433. [\[Cross Ref\]](#)
- Murashige, T. and F. Skoog 1962. A revised medium for rapid growth and bio assays with tobacco tissue culture. *Plant Physiol.*, **15**: 473-497. [\[Cross Ref\]](#)
- Nino-Liu D., Ronald P. and Bogdanove A. 2006. *Xanthomonas oryzae* pathovars: model pathogens of a model crop. *Mol. Plant Pathol.*, **7**: 303–324. [\[Cross Ref\]](#)

- Nithya S., Kumam, Y., Varanavasiappan, S., Kumar, K.K., Muthamilan, M and Sudhakar, D 2020. Targeted mutation in *elf4G* gene in rice. *Electronic Journal of Plant Breeding*, **11**: 1194-1199.
- Oliva, R., Ji, C. and Atienza-Grande, G. 2019. Broad-spectrum resistance to bacterial blight in rice using genome editing. *Nat. Biotechnol.*, **37**: 1344–1350. [\[Cross Ref\]](#)
- Porebski, S., Bailey, L. G. and Baum, B. R. 1997. Modification of a CTAB DNA extraction protocol for plants containing high polysaccharide and polyphenol components. *Plant Mol. Biol. Rep.*, **15**: 8-15. [\[Cross Ref\]](#)
- Rao, P.S. and Kauffman, H.E. 1977. Potential yield losses in dwarf rice varieties due to bacterial leaf blight in *Indian Phytopathol.*, **90**: 281–284. [\[Cross Ref\]](#)
- Reddy, A.P.K., Saxena, N.P. and Reddy, A.V.R., 1978. Analysis of loss in yield due to incidence of bacterial blight disease of rice. *Indian Phytopathol.*, **31**: 444–447.
- Ricroch, A., Clairand, P. and Harwood, W. 2017. Use of CRISPR systems in plant genome editing: toward new opportunities in agriculture. *Emerging Top. Life Sci.*, **1**: 169-182. [\[Cross Ref\]](#)
- Sheoran, O.P., Tonk, D.S., Kaushik, L.S., Hasija, R.C. and Pannu, R.S. 1998. Statistical software package for agricultural research workers.
- Shweta S., Varnavasiappan, S., Kumar K. K., Sudhakar, D., Arul, L. and Kokiladevi, E. 2020. Protocol optimization for rapid and efficient callus induction and *in-vitro* regeneration in rice (*Oryza sativa* L.) cv. CO 51. *Electronic Journal of Plant Breeding*, **11**: 755-759.
- Sun, Y., Zhang, X., Wu, C., He, Y., Ma, Y., Hou, H. and Xia, L. 2016. Engineering herbicide-resistant rice plants through CRISPR/Cas9-mediated homologous recombination of acetolactate synthase. *Mol. Plant.*, **9**: 628-631. [\[Cross Ref\]](#)
- Xie, K., Minkenberg, B. and Yang, Y. 2015. Boosting CRISPR/Cas9 multiplex editing capability with the endogenous tRNA-processing system. *PNAS*, **112**: 3570-5. [\[Cross Ref\]](#)
- Xie, X., Ma, X., Zhu, Q., Zeng, D., Li, G. and Liu Y-G. 2017. CRISPR-GE: A Convenient Software Toolkit for CRISPR-Based Genome Editing. *Mol Plant.*, **10**: 1246-1249. [\[Cross Ref\]](#)
- Yinggen K, Hui, S. and Yuan, M. 2017. *Xanthomonas oryzae* pv. *oryzae* Inoculation and growth rate on rice by leaf clipping method, *Bio-protocol.*, **7**: 2568. [\[Cross Ref\]](#)
- Zafar, K., Khan, M. Z., Amin, I., Mukhtar, Z., Yasmin, S., Arif, M., Ejaz, K. and Mansoor, S. 2020. Precise CRISPR-Cas9 mediated genome editing in super basmati rice for resistance against bacterial blight by targeting the major susceptibility gene. *Front. Plant Sci.*, **11**: 575. [\[Cross Ref\]](#)
- Zhang, H., Zhang, J., Wei, P., Zhang, B., Gou, F., Feng, Z., Mao, Y., Yang, L., Zhang, H., Xu, N. and Zhu, J.K. 2014. The CRISPR/Cas9 system produces specific and homozygous targeted gene editing in rice in one generation. *Plant Biotechnol. J.*, **6**: 797-807. [\[Cross Ref\]](#)